

Literatur zum Artikel

Das Watch-and-Wait-Konzept nach Radiochemotherapie des Rektumkarzinoms – bereits Standard?

- Rouleau-Fournier F, Brown CJ (2019) Can less be more? Organ preservation strategies in the management of rectal cancer. *Curr Oncol* 26: S16–S23
- Krebsregisterdaten, Zf. 2022 https://www.krebsdaten.de/Krebs/EN/Content/Cancer_sites/cancer_sites_node.html
- Sauer R, Becker H, Hohenberger W, et al (2004) Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 351: 1731–1740
- Rödel C, Graeven U, Fietkau R, et al (2015) Oxaliplatin added to fluorouracil-based preoperative chemoradiotherapy and postoperative chemotherapy of locally advanced rectal cancer (the German CAO/ARO/AIO-04 study): final results of the multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 16: 979–989
- Bahadoer RR, Dijkstra EA, van Etten B, et al (2021) Short-course radiotherapy followed by chemotherapy before total mesorectal excision (TME) versus preoperative chemoradiotherapy, TME, and optional adjuvant chemotherapy in locally advanced rectal cancer (RAPIDO): a randomised, open-label, phase 3 trial. *Lancet Oncol* 22: 29–42
- Conroy T, Bosset JF, Etienne PL, et al (2021) Neoadjuvant chemotherapy with FOLFIRINOX and preoperative chemoradiotherapy for patients with locally advanced rectal cancer (UNICANCER-PRODIGE 23): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 22: 702–715
- Fokas E, Schlenska-Lange A, Polat B, et al (2021) Chemoradiotherapy plus induction or consolidation chemotherapy as total neoadjuvant therapy for patients with locally advanced rectal cancer: long-term results of the CAO/ARO/AIO-12 randomized clinical trial. *JAMA Oncol* 8: e215445
- Bujko K, Nowacki MP, Nasierowska-Guttmejer A, et al (2006) Long-term results of a randomized trial comparing preoperative short-course radiotherapy with preoperative conventionally fractionated chemoradiation for rectal cancer. *Br J Surg* 93: 1215–1223
- Sebag-Montefiore D, Stephens RJ, Steele R, et al (2009) Preoperative radiotherapy versus selective postoperative chemoradiotherapy in patients with rectal cancer (MRC CR07 and NCIC-CTG C016): a multicentre, randomised trial. *Lancet* 373: 811–820
- Bregendahl S, Emmertsen KJ, Lous J, Laurberg S (2013) Bowel dysfunction after low anterior resection with and without neoadjuvant therapy for rectal cancer: a population-based cross-sectional study. *Colorectal Dis* 15: 1130–1139
- Habr-Gama A, de Souza PM, Ribeiro U, et al (1998) Low rectal cancer: impact of radiation and chemotherapy on surgical treatment. *Dis Colon Rectum* 41: 1087–1096
- Habr-Gama A, Oliva Perez R, Nadalin W, et al (2004) Operative versus nonoperative treatment for stage 0 distal rectal cancer following chemoradiation therapy: long-term results. *Ann Surg* 240: 711–717
- Habr-Gama A, Oliva Perez R, Proscurshim I, et al (2006) Patterns of failure and survival for nonoperative treatment of stage c0 distal rectal cancer following neoadjuvant chemoradiation therapy. *J Gastrointest Surg* 10: 1319–1328
- Habr-Gama A, Pagin São Julião G, Borba Vailati B, et al (2019) Organ preservation in cT2N0 rectal cancer after neoadjuvant chemoradiation therapy: the impact of radiation therapy dose-escalation and consolidation chemotherapy. *Ann Surg* 269: 102–107
- Dossa F, Chesney TR, Acuna SA, et al (2017) A watch-and-wait approach for locally advanced rectal cancer after a clinical complete response following neoadjuvant chemoradiation: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2: 501–513
- van der Valk MJM, Hilling DE, Bastiaannet E, et al (2018) Long-term outcomes of clinical complete responders after neoadjuvant treatment for rectal cancer in the International Watch & Wait Database (IWWD): an international multicentre registry study. *Lancet* 391: 2537–2545
- Fernandez LM, et al (2021) Conditional recurrence-free survival of clinical complete responders managed by watch and wait after neoadjuvant chemoradiotherapy for rectal cancer in the International watch & wait database: a retrospective, international, multicentre registry study. *Lancet Oncol* 22: 43–50
- Dattani M, Heald RJ, Goussous G, et al (2018) Oncological and survival outcomes in watch and wait patients with a clinical complete response after neoadjuvant chemoradiotherapy for rectal cancer: a systematic review and pooled analysis. *Ann Surg* 268: 955–967
- Maas M, et al (2015) Assessment of clinical complete response after chemoradiation for rectal cancer with digital rectal examination, endoscopy, and MRI: selection for organ-saving treatment. *Ann Surg Oncol* 22: 3873–3880
- Nasir I, et al (2019) Salvage surgery for local regrowths in watch & wait – are we harming our patients by deferring the surgery? *Eur J Surg Oncol* 45: 1559–1566
- Habr-Gama A, Pagin São Julião G, Gama-Rodrigues J, et al (2017) Baseline T classification predicts early tumor regrowth after nonoperative management in distal rectal cancer after extended neoadjuvant chemoradiation and initial complete clinical response. *Dis Colon Rectum* 60: 586–594
- Lopez-Campos F, et al (2020) Watch and wait approach in rectal cancer: current controversies and future directions. *World J Gastroenterol* 26: 4218–4239
- On J, Shim J, Aly EH (2019) Systematic review and meta-analysis on outcomes of salvage therapy in patients with tumour recurrence during 'watch and wait' in rectal cancer. *Ann R Coll Surg Engl* 101: 441–452
- Nagtegaal ID, Glynne-Jones R (2020) How to measure tumour response in rectal cancer? An explanation of discrepancies and suggestions for improvement. *Cancer Treat Rev* 84: 101964
- Boldrini L, et al (2019) Delta radiomics for rectal cancer response prediction with hybrid 0.35 T magnetic resonance-guided radiotherapy (MRgRT): a hypothesis-generating study for an innovative personalized medicine approach. *Radiol Med* 124: 145–153
- ACO/ARO/AIO. Konsentierte Stellungnahme der ACO, der AIO und der ARO zum „watch and wait“-Konzept mit intendiertem Organerhalt bei Rektumkarzinomen des mittleren und unteren Drittels. http://aco-chirurgie.de/wp-content/uploads/2020/08/ACO-Rektum-ASCO-2020-ww-und-Organerhalt_RR_CR_RH-Endfassung-2020-08-17.pdf. 2020 [19.12.2021]