

Literatur zum Artikel

Minimalinvasive Fetalchirurgie bei Zwerchfellhernie und Spina bifida

1. Carpenter R Jr, Strasburger JF, Garson A Jr (1986) Fetal ventricular pacing for hydrops secondary to complete atrioventricular block. *J Am Coll Cardiol* 8: 1434–1436
2. Allan LD, Maxwell DJ, Carminati M, Tynan MJ (1995) Survival after fetal aortic balloon valvoplasty. *Ultrasound Obstet Gynecol* 5: 90–91
3. Bruner JP, Tulipan NE, Richards WO (1997) Endoscopic coverage of fetal open myelomeningocele in utero. *Am J Obstet Gynecol* 176(1 Pt 1): 256–257
4. VanderWall KJ, Skarsgard ED, Filly RA, et al (1997) Fetendo-clip: a fetal endoscopic tracheal clip procedure in a human fetus. *J Pediatr Surg* 32: 970–972
5. Quintero RA, Morales JW, Bornick PW, et al (2000) Minimally invasive intraluminal tracheal occlusion in a human fetus with left congenital diaphragmatic hernia at 27 weeks gestation via direct fetal laryngoscopy. *Prenat Neonat Med* 5: 134–140
6. Deprest J, Gratacos E, Nicolaides KH (2004) Fetoscopic tracheal occlusion (FETO) for severe congenital diaphragmatic hernia: evolution of a technique and preliminary results. *Ultrasound Obstet Gynecol* 24: 121–126
7. Kohl T, Gembruch U, Filsinger B, et al (2006) Early clinical experience with deliberately delayed temporary fetoscopic tracheal occlusion for the prenatal treatment of life-threatening right and left congenital diaphragmatic hernias. *Fetal Diagn Ther* 21: 314–318
8. Ruano R, Yoshisaki CT, da Silva MM, et al (2012) A randomized controlled trial of fetal endoscopic tracheal occlusion versus postnatal management of severe isolated congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 39: 20–27
9. Schaible T, Kohl T, Reinshagen K, et al (2012) Right- versus left-sided congenital diaphragmatic hernia: postnatal outcome at a specialized tertiary care center. *Pediatr Crit Care Med* 13: 66–71
10. Heffez DS, Aryanpur J, Hutchins GM, Freeman JM (1990) The paralysis associated with myelomeningocele: clinical and experimental data implicating a preventable spinal cord injury. *Neurosurgery* 26: 987–992
11. Nicolaides KH, Campbell S, Gabbe SG, Guidetti R (1986) Ultrasound screening for spina bifida: cranial and cerebral signs. *Lancet* 2(8498): 72–74
12. Bahlmann F, Reinhard I, Schramm T, et al (2015) Cranial and cerebral signs in the diagnosis of spina bifida between 18 and 22 weeks of gestation: a German multicenter study. *Prenat Diagn* 35: 228–235
13. Chen FC, Gerhardt J, Entezami M, et al (2017) Detection of spina bifida by first trimester screening – results of the prospective multicenter Berlin IT-study. *Ultraschall Med* 38: 151–157
14. Adzick NS, Thom EA, Spong CY, et al; MOMS Investigators (2011) A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med* 364: 993–1004
15. Kohl T, Tchatcheva K, Weinbach J, et al (2010) Partial amniotic carbon dioxide insufflation (PACI) during minimally invasive fetoscopic surgery: early clinical experience in humans. *Surg Endosc* 24: 432–444
16. Kohl T, Große Hartlage M, Kienitz D, et al (2003) Percutaneous fetoscopic patch coverage of experimental lumbosacral full-thickness skin lesions in sheep – a minimally invasive technique aimed at minimizing maternal trauma from fetal surgery for myelomeningocele. *Surg Endoscopy* 17: 1218–1223
17. Kohl T, Hering R, Heep A, et al (2006) Percutaneous fetoscopic patch coverage of spina bifida aperta in the human – Early clinical experience and potential. *Fetal Diagn Ther* 21: 185–193
18. Kohl T (2014) Percutaneous minimally invasive fetoscopic surgery for spina bifida aperta. Part I: surgical technique and perioperative outcome. *Ultrasound Obstet Gynecol* 44: 515–524
19. Verbeek RJ, Heep A, Maurits NM, et al (2011) Fetal endoscopic myelomeningocele closure preserves segmental neurological function. *Dev Med Child Neurol* 54: 15–22
20. Herrera SR, Leme RJ, Valente PR, et al (2012) Comparison between two surgical techniques for prenatal correction of meningomyelocele in sheep. *Einstein (Sao Paulo)* 10: 455–461
21. Verbeek R, Pastuszka A, Koszutski T, et al (2019) The effect of fetally open, fetoscopic and postnatal myelomeningocele closure on neuromuscular outcome in spina bifida aperta. Abstract presented at the Annual Meeting of the European Society of Pediatric Neurosurgeons, Sept 17–21, 2019, Athens, Greece
22. Graf K, Kohl T, Neubauer BA, et al (2015) Percutaneous minimally-invasive fetoscopic surgery for spina bifida aperta. Part III: postnatal neurosurgical interventions in the first year of life. *Ultrasound Obstet Gynecol* 47: 158–161
23. Degenhardt J, Schürg R, Winarno A, et al (2014) Percutaneous minimal-access fetoscopic surgery for spina bifida aperta. Part II: maternal management and outcome. *Ultrasound Obstet Gynecol* 44: 525–531
24. Sacco A, Van der Veeken L, Bagshaw E, et al (2019) Maternal complications following open and fetoscopic fetal surgery: a systematic review and meta-analysis. *Prenat Diagn* 39: 251–268
25. Degenhardt J, Axt-Fliedner R, Enzensberger C, et al (2014) Peri- and postoperative management for minimally invasive fetoscopic surgery of spina bifida. *Z Geburtshilfe Neonatol* 218: 244–247
26. Belfort MA, Whitehead WE, Shamsirsaz AA, et al (2017) Fetoscopic open neural tube defect repair: development and refinement of a two-port, carbon dioxide insufflation technique. *Obstet Gynecol* 129: 734–743