

Literatur zum Artikel

Im Fokus: Die schmerzhafte Achillessehne

- Hulin BT, Gabbett TJ, Caputi P, et al (2016) Low chronic workload and the acute:chronic workload ratio are more predictive of injury than between-match recovery time: a two-season prospective cohort study in elite rugby league players. *Br J Sports Med* 50: 1009–1012
- Knobloch K, Schreibmüller L, Meller R, et al (2008) Superior Achilles tendon microcirculation in tendinopathy among symptomatic female versus male patients. *Am J Sports Med* 36: 509–514
- Knobloch K, Schreibmüller L, Kraemer R, et al (2010) Gender and eccentric training in Achilles mid-portion tendinopathy. *Knee Surg Sports Traumatol Arthrosc* 18: 648–655
- Hay M, Patricios J, Collins R, et al (2013) Association of type XI collagen genes with chronic Achilles tendinopathy in independent populations from South Africa and Australia. *Br J Sports Med* 47: 569–574
- El Khoury L, Posthumus M, Collins M, et al (2013) Polymorphic variation within the ADAMTS2, ADAMTS14, ADAMTS5, ADAM12 and TIMP2 genes and the risk of Achilles tendon pathology: a genetic association study. *J Sci Med Sport* 16: 493–498
- Hajdu SD, Omoumi P, Baalbaki R, et al (2016) Dual-energy computed tomography-based molecular imaging of cholesterol deposits in Achilles tendon xanthomatosis. *Arthritis Rheumatol* 68: 1687
- Gaida JE, Alfredson L, Kiss ZS, et al (2009) Dyslipidemia in Achilles tendinopathy is characteristic of insulin resistance. *Med Sci Sports Exerc* 41: 1194–1197
- Gaida JE, Alfredson H, Kiss ZS, et al (2010) Asymptomatic Achilles tendon pathology is associated with a central fat distribution in men and a peripheral fat distribution in women: a cross sectional study of 298 individuals. *BMC Musculoskelet Disord* 11: 41
- Gaida JE, Ashe MC, Bass SL, Cook JL (2009) Is adiposity an under-recognized risk factor for tendinopathy? A systematic review. *Arthritis Rheum* 61: 840–849
- Tilley BJ, Cook JL, Docking SI, Gaida JE (2016) Is higher serum cholesterol associated with altered tendon structure or tendon pain? A systematic review. *Br J Sports Med* 0:2016–095965v1
- Robinson JM, Cook JL, Purdam C, et al (2001) The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 35: 335–341
- Lohrer H, Nauck T (2009) Cross-cultural adaptation and validation of the VISA-A questionnaire for German-speaking Achilles tendinopathy patients. *BMC Musculoskelet Disord* 10: 134
- Pingel J, Lu Y, Starborg T, et al (2014) 3D-ultrastructure and collagen composition of healthy and overloaded human tendon: evidence of tenocyte and matrix buckling. *J Anat* 224: 548–555
- Knobloch K, Kraemer R, Lichtenberg A, et al (2006) Achilles tendon and paratendon microcirculation in midportion and insertional tendinopathy in athletes. *Am J Sports Med* 34: 92–97
- Knobloch K (2007) Eccentric training in Achilles tendinopathy: is it harmful to tendon microcirculation? *Br J Sports Med* 41: e2
- Knobloch K, Schreibmüller L, Meller R, et al (2008) Superior Achilles tendon microcirculation in tendinopathy among symptomatic female versus male patients. *Am J Sports Med* 36: 509–514
- Hirschmüller A, Frey V, Konstantinidis L, et al (2012) Prognostic value of Achilles tendon Doppler sonography in asymptomatic runners. *Med Sci Sports Exerc* 44: 199–205
- Andersson G, Danielson P, Alfredson H, Forsgren S (2007) Nerve-related characteristics of ventral paratendinous tissue in chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 15: 1272–1279
- Andersson G, Backman LJ, Christensen J, Alfredson H (2016) Nerve distribution in insertional Achilles tendinopathy – a comparison of bone, bursae and tendon. *Histol Histopathol* 2016; Jun 20.
- Andersson G, Danielson P, Alfredson H, Forsgren S (2009) Presence of substance P and the neurokinin-1 receptor in tenocytes of the human Achilles tendon. *Regul Pept* 150: 81–87
- Scott A, Alfredson H, Forsgren S (2008) VGLuT2 expression in painful Achilles and patellar tendinosis: evidence of local glutamate release by tenocytes. *J Orthop Res* 26: 685–692
- Bjur D, Danielson P, Alfredson H, Forsgren S (2008) Immunohistochemical and in situ hybridization observations favor a local catecholamine production in the human Achilles tendon. *Histo Histopathol* 23: 197–208
- Forsgren S, Alfredson H, Andersson G (2015) Further proof of the existence of a non-neuronal cholinergic system in the human Achilles tendon: Presence of the AchRalpha7 receptor in tendon cells and cells in the peritendinous tissue. *Int Immunopharmacol* 29: 195–200
- Gaida JE, Alfredson H, Forsgren S, Cook JL (2016) A pilot study on biomarkers for tendinopathy: lower levels of serum TNF-alpha and other cytokines in females but not males with Achilles tendinopathy. *BMC Sports Sci Med Rehabil* 8: 5
- Tallon C, Maffulli N, Ewen SW (2001) Ruptured Achilles tendons are significantly more degenerated than tendinopathic tendons. *Med Sci Sports Exerc* 33: 1983–1990
- Intziogianni K, Cassel M, Rauf S, et al (2016) Influence of age and pathology on Achilles tendon properties during a single-leg jump. *Int J Sports Med* 2016 Aug 8.
- Zhang Q, Cai Y, Hua Y, et al (2016) Sonoelastography shows that Achilles tendons with insertional tendinopathy are harder than asymptomatic tendons. *Knee Surg Sports Traumatol Arthrosc* 2016 Jun 24.
- Andersson G, Backman LJ, Christensen J, et al (2016) Nerve distributions in insertional Achilles tendinopathy – a comparison of bone, bursae and tendon. *Histol Histopathol*: 11790
- Dirrichs T, Quack V, Gatz M, et al (2016) Shear wave elastography (SWE) for the evaluation of patients with tendinopathies. *Acad Radiol* 2016 Jun 15
- Zhang Q, Cai Y, Hua Y, et al (2016) Sonoelastography shows that Achilles tendons with insertional tendinopathy are harder than asymptomatic tendons. *Knee Surg Sports Traumatol Arthrosc* [epub ahead of print]
- Knobloch K (2007) Eccentric training in Achilles tendinopathy: is it harmful to tendon microcirculation? *Br J Sports Med* 41:e2
- Langberg H, Ellingsgaard H, Madsen T, et al (2007) Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports* 17: 61–66
- Heinemeier KM, Olesen JL, Haddad F, et al (2007) Expression of collagen and related growth factors in rat tendon and skeletal muscle in response to specific contraction types. *J Physiol* 582: 1303–1316
- Farris DJ, Buckeridge E, Trewartha G, McGuigan MP (2012) The effects of orthotic heel lifts on Achilles tendon force and strain during running. *J Appl Biomech* 28: 511–519
- Wulf M, Wearing SC, Hooper SL, et al (2016) The effect of an in-shoe orthotic heel lift on loading of the Achilles tendon during shoe walking. *J Orthop Sports Phys Ther* 46: 79–86
- Weinert-Aplin RA, Bull AM, McGregor AH (2016) Orthotic heel wedges do not alter hindfoot kinematics and Achilles tendon force during level and inclined walking in healthy individuals. *J Appl Biomech* 32: 160–170
- Scott LA, Munteanu SE, Menz HB (2015) Effectiveness of orthotic devices in the treatment of Achilles tendinopathy: a systematic review. *Sports Med* 45: 95–110
- Mousavizadeh R, Backman L, McCormack RG, Scott A (2015) Dexamethasone decreases substance P expression in human tendon cells: an in vitro study. *Rheumatology (Oxford)* 54: 318–323
- Turmo-Garuz A, Rodas G, Balias R, et al (2014) Can local corticosteroid injection in the retrocalcaneal bursa lead to rupture of the Achilles tendon and the medial head of the gastrocnemius muscle? *Musculoskelet Surg* 98: 121–126
- Muto T, Kokubu T, Mifune Y, et al (2014) Temporary inductions of matrix metalloproteinase-3 (MMP-3) expression and cell apoptosis are associated with tendon degeneration or rupture after corticosteroid injection. *J Orthop Res* 32: 1297–1304

41. Taneja AK, Santos DC (2014) Steroid-induced Kager's fat pad atrophy. *Skeletal Radiol* 43: 1161–1164
42. Coombes BK, Bisset L, Vicenzino B (2010) Efficacy and safety of corticosteroid injections and other injections for management of tendinopathy: a systematic review of randomized controlled trials. *Lancet* 376 (9754): 1751–1767
43. Alfredson H, Ohberg L, Zeisig E, Lorentzon R (2007) Treatment of midportion Achilles tendinosis: similar clinical results with US and CD-guided surgery outside the tendon and sclerosing polidocanol injections. *Knee Surg Sports Traumatol Arthrosc* 15: 1504–1509
44. Willberg L, Sunding K, Ohberg L, et al (2008) Sclerosing injections to treat midportion Achilles tendinosis: a randomised controlled study evaluating two different concentrations of Polidocanol. *Knee Surg Sports Traumatol Arthrosc* 16: 859–864
45. Lind B, Ohberg L, Alfredson H (2008) Sclerosing polidocanol injections in mid-portion Achilles tendinosis: remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 14: 1327–1332
46. Osti L, Buda M, Buono AD, et al (2015) Clinical evidence in the treatment of rotator cuff tears with hyaluronic acid. *Muscles Ligaments Tendons J* 5: 270–275
47. Frizziero A, Salamanna F, Giavaresi G, et al (2015) Hyaluronic acid injections protect patellar tendon from detraining-associated damage. *Histol Histopathol* 30: 1079–1088
48. Kaux JF, Samson A, Crielaard JM (2015) Hyaluronic acid and tendon lesions. *Muscles Ligaments Tendons J* 5: 264–269
49. Kumai T, Muneta T, Tsuchiya A, et al (2014) The short-term effect after a single injection of high-molecular-weight hyaluronic acid in patients with enthesiopathies (lateral epicondylitis, patellar tendinopathy, insertional Achilles tendinopathy, and plantar fasciitis): a preliminary study. *J Orthop Sci* 19: 603–611
50. De Vos RJ, Weir A, van Schie HAT, et al (2010) Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 303: 144–149
51. Krogh TP, Ellingsen T, Christensen R, et al (2016) Ultrasound-guided injection therapy of Achilles tendinopathy with platelet-rich plasma or saline: A randomized, blinded, placebo-controlled trial. *Am J Sports Med* 44: 1990–1997
52. Gerdemeyer L, Mittermayr R, Fuerst M, et al (2015) Current evidence of ESWT in chronic Achilles tendinopathy. *Int J Surg* 24 (Pt B): 154–159
53. Mani-Babu S, Morrissey D, Waugh C, et al (2015) The effectiveness of ESWT in lower limb tendinopathy: a systematic review. *Am J Sports Med* 43: 752–761
54. Rompe JD, Furla J, Maffulli N (2009) Eccentric loading versus eccentric loading plus shock-wave treatment for midportion Achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 37: 463–470
55. Rompe JD, Furla J, Maffulli N (2008) Eccentric loading compared with shock-wave treatment for chronic insertional Achilles tendinopathy: a randomized, controlled trial. *J Bone Joint Surg Am* 90: 52–61
56. Leone L, Raffa S, Vetrano M, et al (2016) Extracorporeal shock wave treatment (ESWT) enhances the in vitro-induced differentiation of human tendon-derived stem/progenitor cells (hTSPCs). *Oncotarget* 7: 6410–6423
57. Ng GY, Chung PY (2012) Effects of a therapeutic laser and passive stretching program for treating tendon overuse. *Photomed Laser Surg* 30: 155–159
58. Mardh A, Lund I (2016) High power Laser for treatment of Achilles tendinosis – a single blind randomized placebo controlled clinical study. *J Lasers Med Sci* 7: 92–98
59. Torres-Silva R, Lopes-Martins RA, Bjordal JM, et al (2015) The low level laser therapy (LLLT) operating in 660nm reduce gene expression of inflammatory mediators in the experimental model of collagenase-induced rat tendinitis. *Laser Med Sci* 30: 1985–1990
60. Chen MH, Huang YC, Sun JS, et al (2015) Second messengers mediating the proliferation and collagen synthesis of tenocytes induced by low-level laser irradiation. *Lasers Med Sci* 30: 263–272
61. Guerra Fda R, Vieira CP, Almeida MS, et al (2013) LLLT improves tendon healing through increase of MMP activity and collagen synthesis. *Lasers Med Sci* 28: 1281–1288
62. Ott OJ, Jeremias C, Gajpl US, et al (2015) Radiotherapy for benign allodynia. Long-term results of the Erlangen dose optimization trial. *Strahlenther Onkol* 191: 979–984
63. Lohrer H, David D, Nauck T (2016) Surgical treatment for Achilles tendinopathy – a systematic review. *BMC Musculoskelet Disord* 17: 207
64. Baltes TP, Zwiers R, Wiegerinck JI, van Dijk CN (2016) Surgical treatment for midportion Achilles tendinopathy: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2016 May 12.
65. Alfredson H, Pietilä T, Öhberg L, Lorentzon R (1998) Achilles tendinosis and calf muscle strength: the effect of short-term immobilization after surgical treatment. *Am J Sports Med* 26: 166–171
66. Thermann H, Benetos IS, Panelli C, et al (2009) Endoscopic treatment of chronic mid-portion Achilles tendinopathy: novel technique with short-term results. *Knee Surg Sports Traumatol Arthrosc* 17: 1264–1269
67. Alfredson H (2016) Low recurrence rate after mini surgery outside the tendon combined with short rehabilitation in patients with midportion Achilles tendinopathy. *Open Access J Sports Med* 7: 51–54