

Literatur zum Artikel

Chronischer Schmerz nach Leistenhernienoperation

1. Aasvand EK, Kehlet H (2005) Chronic postoperative pain: the case of inguinal herniorrhaphy. *Br J Anaesth* 95: 69–76
2. Aasvang EK, Gmaehle E, Hansen JB, et al (2010) Predictive risk factors for persistent postherniotomy pain. *Anesthesiology* 112: 957–969
3. Alfieri S, Amid PK, Campanelli G, et al (2011) International guidelines for prevention and management of post-operative chronic pain following inguinal hernia surgery. *Hernia* 15: 239–249
4. Alfieri S, Rotondi F, Di Miceli D, et al (2006) Chronic pain after inguinal hernia mesh repair: possible role of surgical manipulation of the inguinal nerves. A prospective multicentre study of 973 cases. *Chir Ital* 58: 23–31
5. Apkarian AV, Baliki MN, Geha PY (2009) Towards a theory of chronic pain. *Prog Neurobiol* 87: 81–97
6. Arlt GD, Huhn U, Kersten CC (2010) Surgery for chronic inguinal pain: neurectomy, mesh explantation, or both? In: Schumpelick V, Fitzgibbons RJ (eds) *Hernia repair sequelae*. Springer, Berlin. S 294–298
7. Arlt GD, Dübbers S, Huhn U (2014) Long-term outcome of surgery for chronic pain following inguinal hernia repair. 36th International Congress of the European Hernia Society, Edinburgh 29.–31.5.2014. *Hernia* 18 (Suppl 2): P23
8. Aroori S, Spence RA (2007) Chronic pain after hernia surgery – an informed consent issue. *Ulster Med J* 76: 136–140
9. Bay-Nielsen M, Nilsson E, Nordin P, Kehlet H (2004) Chronic pain after open mesh and sutured repair of indirect inguinal hernia in young males. *Br J Surg* 91: 1373–1376
10. Bischoff JM, Ringsted TK, Petersen M, et al (2014) A capsaicin (8%) patch in the treatment of severe persistent inguinal postherniorrhaphy pain: a randomized, double-blind, placebo-controlled trial. *PLoS One* 9(10): e109144
11. Campanelli G, Pascual MH, Hoferlin A, et al (2012) Randomized, controlled, blinded trial of Tisseel/Tissucol for mesh fixation in patients undergoing Lichtenstein technique for primary inguinal repair: results of the TIMELI trial. *Ann Surg* 255: 650–657
12. Courtney CA, Duffy K, Serpell MG, O'Dwyer PJ (2002) Outcome of patients with severe chronic pain following repair of groin hernia. *Br J Surg* 89: 1310–1314
13. Delikoukos S, Fafoulakis F, Christodoulidis G, et al (2008) Re-operation due to severe late-onset persisting groin pain following anterior inguinal hernia repair with mesh. *Hernia* 12: 593–595
14. Demirer S, Kepenekci I, Evirgen O, et al (2006) The effect of polypropylene mesh on ilioinguinal nerve in open mesh repair of groin hernia. *J Surg Res* 13: 175–181
15. Dickinson KJ, Thomas M, Fawole AS, et al (2008) Predicting chronic post-operative pain following laparoscopic inguinal hernia repair. *Hernia* 12: 597–601
16. Dominguez CA, Kalliomäki ML, Gunnarsson U, et al (2013) The DQB1*03:02 HLA haplotype is associated with increased risk of chronic pain after inguinal hernia surgery and lumbar disc herniation. *Pain* 154: 427–433
17. Ece I, Yilmaz H (2018) An overlooked complication of the inguinal hernia repair: dysejaculation. *Turk J Surg* 34: 1–4
18. Eker HH, Langeveld HR, Klitsie PJ, et al (2012) Randomized clinical trial of total extraperitoneal inguinal hernioplasty vs. Lichtenstein repair: a long-term follow-up study. *Arch Surg* 147: 256–260
19. Fränneby U, Sandblom G, Nordin P, et al (2006) Risk factors for long-term pain after hernia surgery. *Ann Surg* 244: 212–219
20. Gupta A, Mayer EA, Fling C, et al (2017) Sex-based differences in brain alterations across chronic pain conditions. *J Neurosci Res* 95: 604–616
21. Gutlic N, Petersson U, Rogmark P, Montgomery A (2018) The relevance of sexual dysfunction related to groin pain after inguinal hernia repair – the sexIHQ short form questionnaire assessment. *Front Surg* 5: 15
22. Johner A, Faulds J, Wiseman SM (2011) Planned ilioinguinal nerve excision for prevention of chronic pain after inguinal hernia repair: a meta-analysis. *Surgery* 150: 534–541
23. Joshi GP, Rawal N, Kehlet H, et al (2012) Evidence-based management of post-operative pain in adults undergoing open inguinal hernia surgery. *Br J Surg* 99: 168–185
24. Kalliomäki ML, Meyerson J, Gunnarsson U, et al (2008) Long-term pain after inguinal hernia repair in a population-based cohort: risk factors and interference with daily activities. *Eur J Pain* 12: 214–225
25. Kalliomäki ML, Sandblom G, Gunnarsson U, Gordh T (2009) Persistent pain after groin hernia surgery: a qualitative analysis of pain and its consequences for quality of life. *Acta Anaesthesiol Scand* 53: 236–246
26. Karampinis I, Weiss J, Pilz L, et al (2017) Transabdominal laparoscopic retroperitoneal neurectomy for chronic pain after inguinal hernia repair and appendectomy – a matched-pair study. *BMC* 17: 85
27. Lange JFM, Wijsmuller AR, van Geldere D, et al (2009) Feasibility study of three-nerve-recognizing Lichtenstein procedure for inguinal hernia. *Br J Surg* 96: 1210–1214
28. Loos MJA, Scheltinga MRM, Roumen RMH (2010) Tailored neurectomy for treatment of postherniorrhaphy inguinal neuralgia. *Surgery* 147: 275–281
29. Lundström KJ, Holmberg H, Montgomery A, Nordin P (2018) Patient-reported rates of chronic pain and recurrence after groin hernia repair. *Br J Surg* 105: 106–112
30. Moore AM, Bjurström MF, Hiatt JR, et al (2016) Efficacy of retroperitoneal triple neurectomy for refractory neuropathic inguino-dynia. *Am J Surg* 212: 1126–1132
31. Öberg S, Andresen K, Rosenberg J (2018) Decreasing prevalence of chronic pain after laparoscopic groin hernia repair: a nationwide cross-sectional questionnaire study. *Surg Today*. doi: 10.1007/s00595-018-1664-5 [Epub ahead of print]
32. Powell R, Johnston M, Smith WC, et al (2012) Psychological risk factors for chronic post-surgical pain after inguinal hernia repair surgery: a prospective cohort study. *Eur J Pain* 16: 600–610
33. Quyn AJ, Weatherhead KM, Daniel T (2012) Chronic pain after open inguinal hernia surgery: suture fixation versus self-adhesive mesh repair. *Langenbecks Arch Surg* 397: 1215–1218
34. Reinhold WM, Nehls J, Eggert A (2011) Nerve management and chronic pain after open inguinal hernia repair: a prospective two phase study. *Ann Surg* 254: 163–168
35. Rozen D, Ahn J (2006) Pulsed radiofrequency for the treatment of ilioinguinal neuralgia after inguinal herniorrhaphy. *Mt Sinai J Med* 73: 716–718
36. Sajid MS, Kalra L, Parampalli U, et al (2013) A systematic review and meta-analysis evaluating the effectiveness of lightweight mesh against heavyweight mesh in influencing the incidence of chronic groin pain following laparoscopic inguinal hernia repair. *Am J Surg* 205: 726–736
37. Sajid MS, Leaver C, Baig MK, Sains P (2012) Systematic review and meta-analysis of the use of lightweight versus heavyweight mesh in open inguinal hernia repair. *Br J Surg* 99: 29–37
38. Schmidt-Wilcke T, Kaitys A, Ichesco E, et al (2014) Changes in clinical pain in fibromyalgia patients correlate with changes in brain activation in the cingulate cortex in a response inhibition task. *Pain Med* 15: 1346–1358
39. Singh AN, Bansal VK, Misra MC, et al (2012) Testicular functions, chronic groin pain, and quality of life after laparoscopic and open mesh repair of inguinal hernia: a prospective randomized controlled trial. *Surg Endosc* 26: 1304–1317
40. Slooter GD, Zwaans WAR, Perquin CW, et al (2018) Laparoscopic mesh removal for otherwise intractable inguinal pain following endoscopic hernia repair is feasible, safe and may be effective in selected patients. *Surg Endosc* 32: 1613–1619
41. Uzzaman MM, Ratnasingham K, Ashraf N (2012) Meta-analysis of randomized controlled trials comparing lightweight and heavyweight mesh for Lichtenstein inguinal hernia repair. *Hernia* 16: 505–518
42. Verhagen T, Loos MJA, Scheltinga MRM, et al (2018) The groin pain trial: a randomized controlled trial of injection therapy versus neurectomy for postherniorrhaphy inguinal neuralgia. *Ann Surg* 267: 841–845
43. Woolf CJ (2011) Central sensitization. Implications for the diagnosis and treatment of pain. *Pain* 152 (3 Suppl): S2–15
44. Zacest AC, Magill ST, Anderson VC, et al (2010) Long-term outcome following ilioinguinal neurectomy for chronic pain. *J Neurosurg* 112: 784–789