

Literatur zum Artikel

Gliedmaßenverlust bei Weichgewebesarkomen

1. Enneking WF (1983) *Musculoskeletal tumor surgery*. Churchill Livingstone, New York
2. Enzinger FM, Weiss SW (1995) In: *Soft tissue tumors*, 3rd ed. Mosby, St. Louis
3. The ESMO/European Sarcoma Network Working Group (2014) Soft tissue and visceral sarcomas: ESMO clinical practice guidelines. *Ann Oncol* 25 (Suppl 3) III102–III112
4. Eggermont AM, Schraffordt Koops H, Klausner JM, et al (1996) Isolated limb perfusion with tumor necrosis factor and melphalan for limb salvage in 186 patients with locally advanced soft tissue extremity sarcomas. The cumulative multicenter European experience. *Ann Surg* 224: 756–764
5. Schwarzbach MHM, Hormann Y, Hinz U, et al (2005) Results of limb-sparing surgery with vascular replacement for soft tissue sarcoma in the lower extremity. *J Vasc Surg* 42: 88–96
6. Schwarzbach MH, Hormann Y, Hinz U (2006) Clinical results of surgery for retroperitoneal sarcoma with major blood vessel involvement. *J Vasc Surg* 44: 46–55
7. Koperna T, Teleky B, Vogl S (1996) Vascular reconstruction for limb salvage in sarcoma of the lower extremity. *Arch Surg* 131: 1103–1107
8. Nabisan RN, Karakousis CP (1987) Vascular reconstruction for limb salvage in soft tissue sarcoma. *Surgery* 101: 668–677
9. Thomas JE, Piepgras DG, Scheithauer B, et al (1983) Neurogenic tumors of the sciatic nerve. A clinicopathologic study of 35 cases. *Mayo Clin Proc* 58: 640–647
10. Bickels J, Wittig JC, Kollender Y, et al (2002) Sciatic nerve resection: is that truly an indication for amputation? *Clin Orthop Relat Res* 399: 201–204
11. Brooks AD, Gold JS, Graham D, et al (2002) Resection of the sciatic, peroneal, or tibial nerves: assessment of functional status. *Ann Surg Oncol* 9: 41–47
12. Fuchs B, Davis AM, Wunder JS, et al (2001) Sciatic nerve resection in the thigh: a functional evaluation. *Clin Orthop Relat Res* 382: 34–41
13. Clarkson PW, Griffin AM, Catton CN, et al (2005) Epineural dissection is a safe technique that facilitates limb salvage surgery. *Clin Orthop Relat Res* 438: 92–96
14. Cardona S, Schwarzbach M, Hinz U, et al (2003) Evaluation of F18-deoxyglucose positron emission tomography (FDG-PET) to assess the nature of neurogenic tumors. *Eur J Surg Oncol* 29: 536–541
15. Sweiti H, Tamimi N, Bormann F, et al (2018) Limb-salvage surgery of soft tissue sarcoma with sciatic nerve involvement. *Sarcoma* (in press)
16. Eggermont AM (2003) Isolated limb perfusion in the management of locally advanced extremity soft tissue sarcoma. *Surg Oncol Clin N Am* 83: 371–384
17. Podleska LE, Schwindinghammer B, Grabellus F, et al (2017) Isolierte Extremitätenperfusionen bei Liposarkomen: Histopathologisches Ansprechen und Subgruppenanalyse nach TNF-Melphalan ILP. *Chirurg* 88: 429–436
18. Jakob J, Hohenberger P (2016) Role of isolated limb perfusion with recombinant Human Tumor Necrosis Factor alpha and Melphalan in locally advanced extremity soft tissue sarcoma. *Cancer* 122: 2624–2632
19. Schwarzbach MHM, Lehnert T, Willeke F, et al (1996) Ergebnisse der isolierten hyperthermen Extremitätenperfusion bei Weichteilsarkomen im Rahmen eines multimodalen Behandlungskonzeptes. *Chirurg* 67: 1237–1243
20. Grabellus F, Kraft C, Sheu-Grabellus SY, et al (2011) Tumor vascularization and histopathologic regression of soft tissue sarcomas treated with isolated limb perfusion with TNF-alpha and melphalan. *J Surg Oncol* 103: 371–379
21. Herbert SH, Corn BW, Solin LJ, et al (1993) Limb-preserving treatment for soft tissue sarcomas of the extremities. The significance of surgical margins. *Cancer* 72: 1230–1238
22. Stojadinovic A, Leung D, Hoos A, et al (2002) Analysis of the prognostic significance of microscopic margins in 2,084 localized primary adult soft tissue sarcomas. *Ann Surg* 235: 424–434
23. Nystrom LM, Reimer NB, Reith JD, et al (2013) Multidisciplinary management of soft tissue sarcoma. *Scient World J* 2013: 852462
24. O'Sullivan B, Davis AM, Turcotte R, et al (2002) Preoperative versus postoperative radiotherapy in soft-tissue sarcoma of the limbs: a randomised trial. *Lancet* 359(9325): 2235–2241
25. Dagan R, Indelicato DJ, McGee L, et al (2012) The significance of a marginal excision after preoperative radiation therapy for soft tissue sarcoma of the extremity. *Cancer* 118: 3199–3207
26. Abatzoglou S, Turcotte RE, Adoubali A, et al (2010) Local recurrence after initial multidisciplinary management of soft tissue sarcoma: is there a way out? *Clin Orthop Relat Res* 468: 3012–3018
27. Novais EN, Demiralp B, Alderete J, et al (2010) Do surgical margin and local recurrence influence survival in soft tissue sarcomas? *Clin Orthop Relat Res* 468: 3003–3011
28. Bonvalot S, Laplanche A, Lejeune F, et al (2005) Limb salvage with isolated perfusion for soft tissue sarcoma: could less TNF-alpha be better? *Ann Oncol* 16: 1061–1068
29. Pennacchioli E, Deraco M, Mariani L, et al (2007) Advanced extremity soft tissue sarcoma: prognostic effect of isolated limb perfusion in a series of 88 patients treated at a single institution. *Ann Surg Oncol* 14: 553–559
30. Rastrelli M, Campana LG, Valpione S, et al (2016) Hyperthermic isolated limb perfusion in local advanced limb soft tissue sarcoma: a 24-year single-centre experience. *Int J Hyperthermia* 32: 165–172
31. Cherix S, Speiser M, Matter M, et al (2008) Isolated limb perfusion with tumor necrosis factor and melphalan for non-resectable soft tissue sarcomas: long-term results on efficacy and limb salvage in a selected group of patients. *J Surg Oncol* 98: 148–155
32. Jakob J, Tunn PU, Hayes AJ, et al (2014) Oncological outcome of primary non-metastatic soft tissue sarcoma treated by neoadjuvant isolated limb perfusion and tumor resection. *J Surg Oncol* 109: 786–790
33. Schwarzbach MHM, Jordan M, Beidatsch D, et al (2017) Isolierte hypertherme Extremitätenperfusion mit rh TNF-alpha und Melphalan: Extremitätenerhalt bei lokal fortgeschrittenen Weichgewebesarkomen. *CHAZ* 18: 179–184
34. Schwarzbach MH, Koesters R, Germann A, et al (2004) Comparable transforming capacities and differential gene expression patterns of variant FUS/CHOP fusion transcripts derived from soft tissue liposarcomas. *Oncogene* 23: 6798–6805
35. Schwarzbach MHM, Hinz U, Dimitrakopoulou-Strauss A, et al (2005) Prognostic significance of preoperative [18-F] fluorodeoxyglucose (FDG) positron emission tomography (PET) imaging in patients with resectable soft tissue sarcomas. *Ann Surg* 241: 286–294
36. Schwarzbach MHM, Dimitrakopoulou-Strauss A, Willeke F, et al (2000) Clinical value of [18-F] fluorodeoxyglucose positron emission tomography imaging in soft tissue sarcomas. *Ann Surg* 231: 380–386
37. Schwarzbach MHM, Dimitrakopoulou-Strauss A, Mechttersheimer G, et al (2001) Prospective assessment of masses suspicious for liposarcomas by fluorine-18-2-fluoro-2-deoxy-D-glucose (FDG) positron emission tomography (PET). *Anticancer Res* 21: 3609–3614
38. Prewitt TW, Alexander HR, Sindelar WF (1995) Hemipelvectomy for soft tissue sarcoma: clinical results in fifty-three patients. *Surg Oncol* 4: 261–269
39. Williard WC, Hajdu SI, Casper ES, Brennan MF (1992) Comparison of amputation with limb-sparing operations for adult soft tissue sarcoma of the extremity. *Ann Surg* 215: 269–275
40. Conrad EU, Springfield D, Peabody TD (1998) Pelvis. In: Simon MA, Springfield D (eds) *Surgery for bone and soft tissue tumors*. Lippincott-Raven, Philadelphia, 323–341
41. Dorsi MJ, Zwagil ZS, Hsu W, Belzberg AJ (2011) Epithelioid sarcoma of the tibial portion of the sciatic nerve. *Clin Neurol Neurosurg* 113: 506–508

42. Davis AM, Devlin M, Griffin AM, et al (1999) Functional outcome in amputation versus limb sparing of patients with lower extremity sarcoma: a matched case-control study. *Arch Phys Med Rehabil* 80: 615–618
43. O'Connor MI (1998) Surgical management of malignant soft tissue tumors. In Simon MA, Springfield D (eds) *Surgery for bone and soft tissue tumors*. Lippincott-Raven, Philadelphia, S 555–565
44. Gundle KR, Kafchinski L, Gupta S, et al (2018) Analysis of the margin classification system for assessing the risk of local recurrence after soft tissue sarcoma resection. *J Clin Oncol* 36: 704–709
45. Schwarzbach MH, et al (2012) Moderne Chirurgie retroperitonealer Weichgewebesarkome: Primärtumore. *CHAZ* 13: 207–215
46. Schwarzbach MH, et al (2012) Moderne Chirurgie retroperitonealer Weichgewebesarkome: Lokalrezidive. *CHAZ* 13: 335–340
47. Schwarzbach MH, et al (2014) Weichgewebesarkome mit Blutgefäßbeteiligung. *CHAZ* 15: 188–195