

Literatur zum Artikel

Reduktion der Narbenhernieninzidenz durch prophylaktische Netzimplantation nach offener Abdominalbehandlung (PROMOAT)

1. Coccolini F, Montori G, Ceresoli M, et al (2017) The role of open abdomen in non-trauma patient: WSES Consensus Paper. *World J Emerg Surg* 12: 39
2. Sartelli M, Abu-Zidan FM, Ansaloni L, et al (2015) The role of the open abdomen procedure in managing severe abdominal sepsis: WSES position paper. *World J Emerg Surg* 10: 35
3. Atema JJ, Gans SL, Boermeester MA (2015) Systematic review and meta-analysis of the open abdomen and temporary abdominal closure techniques in non-trauma patients. *World J Surg* 39: 912–925
4. Borab ZM, Shakir S, Lanni MA, et al (2017) Does prophylactic mesh placement in elective, midline laparotomy reduce the incidence of incisional hernia? A systematic review and meta-analysis. *Surgery* 161: 1149–1163
5. Willms A, Günsen C, Schaaf S, et al (2015) Management of the open abdomen using vacuum-assisted wound closure and mesh-mediated fascial traction. *Langenbecks Arch Surg* 400: 91–99
6. Willms AG, Schaaf S, Zimmermann N, et al (2019) The significance of visceral protection in preventing enteroatmospheric fistulae during open abdomen treatment in patients with secondary peritonitis: a propensity score-matched case-control analysis. *Ann Surg*. doi: 10.1097/SLA.0000000000003440 [ahead of print]
7. Brown SR, Goodfellow PB (2005) Transverse versus midline incisions for abdominal surgery. *Cochrane Database Syst Rev* 4: CD005199
8. Willms A, Schaaf S, Schwab R, et al (2017) Intensive care and health outcomes of open abdominal treatment: long-term results of vacuum-assisted wound closure and mesh-mediated fascial traction (VAWCM). *Langenbecks Arch Surg* 402: 481–492
9. Bjarnason T, Montgomery A, Ekberg O, et al (2013) One-year follow-up after open abdomen therapy with vacuum-assisted wound closure and mesh-mediated fascial traction. *World J Surg* 37: 2031–2038
10. Brandl A, Laimer E, Perathoner A, et al (2014) Incisional hernia rate after open abdomen treatment with negative pressure and delayed primary fascia closure. *Hernia* 18:105–111
11. Willms A, Schaaf S, Schwab R, et al (2016) Abdominal wall integrity after open abdomen: long-term results of vacuum-assisted wound closure and mesh-mediated fascial traction (VAWCM). *Hernia* 20: 849–858
12. Petersson P, Montgomery A, Petersson U (2019) Vacuum-assisted wound closure and permanent onlay mesh-mediated fascial traction: a novel technique for the prevention of incisional hernia after open abdomen therapy including results from a retrospective case series. *Scand J Surg* 108: 216–226
13. López-Cano M, García-Alamino JM, Antoniou SA, et al (2018) EHS clinical guidelines on the management of the abdominal wall in the context of the open or burst abdomen. *Hernia* 22: 921–939
14. Bleier JIS, Resnick AS (2009) Complications of incisional hernia repair. *Semin Colon Rectal Surg* 20: 125–130
15. Azar FK, Crawford TC, Poruk KE, et al (2017) Ventral hernia repair in patients with abdominal loss of domain: an observational study of one institution's experience. *Hernia* 21: 245–252
16. Muysoms FE, Dietz UA (2016) Prophylaktische Netze an der Bauchdecke. *Chirurg* 87: 751–761
17. Berrevoet F (2018) Prevention of incisional hernias after open abdomen treatment. *Front Surg* 5: 11
18. Muysoms FE, Antoniou SA, Bury K, et al (2015) European Hernia Society guidelines on the closure of abdominal wall incisions. *Hernia* 19: 1–24
19. Jairam AP, Timmermans L, Eker HH, et al (2017) Prevention of incisional hernia with prophylactic onlay and sublay mesh reinforcement versus primary suture only in midline laparotomies (PRIMA): 2-year follow-up of a multicentre, double-blind, randomised controlled trial. *Lancet* 390(10094): 567–576
20. Depuydt M, Allaey M, Carvalho LA de, et al (2021) Prophylactic mesh after midline laparotomy: evidence is out there, but why do surgeons hesitate? *World J Surg* 45: 1349–1361
21. Payne R, Aldwinckle J, Ward S (2017) Meta-analysis of randomised trials comparing the use of prophylactic mesh to standard midline closure in the reduction of incisional herniae. *Hernia* 21: 843–853
22. Bueno-Lledó J, Torregrosa-Gallud A, Sala-Hernandez A, et al (2017) Predictors of mesh infection and explantation after abdominal wall hernia repair. *Am J Surg* 213: 50–57
23. National Healthcare Safety Network, Centers for Disease Control and Prevention (2017) Surgical site infection (SSI) event. <http://www.cdc.gov/nhsn/pdfs/pscmanual/9pscscscurrent.pdf>. [2. Juli 2020]
24. Clavien PA, Barkun J, Oliveira ML de, et al (2009) The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg* 250: 187–196
25. Hawker GA, Mian S, Kendzerska T, et al (2011) Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res (Hoboken)* 63: S240-52
26. Schaaf S, Schwab R, Günsen C, et al (2020) Prophylactic onlay mesh implantation during definitive fascial closure after open abdomen therapy (PROMOAT): absorbable or non-absorbable? Methodical description and results of a feasibility study. *Front Surg* 7: 578565
27. Muysoms FE, Detry O, Vierendeels T, et al (2016) Prevention of incisional hernias by prophylactic mesh-augmented reinforcement of midline laparotomies for abdominal aortic aneurysm treatment: a randomized controlled trial. *Ann Surg* 263: 638–645
28. Kohler A, Lavanchy JL, Lenoir U, et al (2019) Effectiveness of prophylactic intraperitoneal mesh implantation for prevention of incisional hernia in patients undergoing open abdominal surgery: a randomized clinical trial. *JAMA Surg* 154: 109–115
29. Indrakusuma R, Jalalzadeh H, van der Meij JE, et al (2018) Prophylactic mesh reinforcement versus sutured closure to prevent incisional hernias after open abdominal aortic aneurysm repair via midline laparotomy: a systematic review and meta-analysis. *Eur J Vasc Endovasc Surg* 56: 120–128
30. Wanhainen A (2018) To mesh or not to mesh? That is the question! *Eur J Vasc Endovasc Surg* 56: 129
31. San Miguel C, Melero D, Jiménez E, et al. (2018) Long-term outcomes after prophylactic use of onlay mesh in midline laparotomy. *Hernia* 22: 1113–1122
32. Glauser PM, Brosi P, Speich B, et al (2019) Prophylactic intraperitoneal onlay mesh following midline laparotomy-long-term results of a randomized controlled trial. *World J Surg* 43: 1669–1675
33. Ahmed J, Hasnain N, Fatima I, et al (2020) Prophylactic mesh placement for the prevention of incisional hernia in high-risk patients after abdominal surgery: a systematic review and meta-analysis. *Cureus* 12: e10491
34. Heger P, Feißt M, Krisam J, et al (2019) Hernia reduction following laparotomy using small stitch abdominal wall closure with and without mesh augmentation (the HULC trial): study protocol for a randomized controlled trial. *Trials* 20: 738
35. Burns FA, Heywood EG, Challand CP, et al (2020) Is there a role for prophylactic mesh in abdominal wall closure after emergency laparotomy? A systematic review and meta-analysis. *Hernia* 24: 441–447
36. Jakob MO, Schwarz C, Haltmeier T, et al (2018) Mesh-augmented versus direct abdominal closure in patients undergoing open abdomen treatment. *Hernia* 22: 785–792
37. Tansawat A, Numthavaj P, Techapongsatorn S, et al (2020) Mesh position for hernia prophylaxis after midline laparotomy: A systematic review and network meta-analysis of randomized clinical trials. *Int J Surg* 83: 144–151