

Literatur zum Artikel

Die posttraumatische Gonarthrose

1. Pickering RD (1984) Posttraumatic arthritis. *Can Fam Physician* 30: 1511–1513
2. Dixel J, Beyer F, Lützner C, et al (2016) TKA for posttraumatic osteoarthritis is more complex and needs more surgical resources. *Orthopedics* 39 (3 Suppl): S36–40
3. Furman BD, Mangiapani DS, Zeitler E, et al (2014) Targeting pro-inflammatory cytokines following joint injury: acute intra-articular inhibition of interleukin-1 following knee injury prevents post-traumatic arthritis. *Arthritis Res Ther* 16: R134
4. Houdek MT, Watts CD, Shannon SF, et al (2016) Posttraumatic total knee arthroplasty continues to have worse outcome than total knee arthroplasty for osteoarthritis. *J Arthroplasty* 31: 118–123
5. Kester BS, Minhas SV, Vigdorich JM, Schwarzkopf R (2016) Total knee arthroplasty for posttraumatic osteoarthritis: is it time for a new classification? *J Arthroplasty* 31: 1649–1653
6. Wu LD, Xiong Y, Yan SG, Yang QS (2005) Total knee replacement for posttraumatic degenerative arthritis of the knee. *Chin J Traumatol* 8: 195–199
7. Chun KC, Kim KM, Chun CH (2013) Infection following total knee arthroplasty. *Knee Surg Relat Res* 25: 93–99
8. Dinneen A, Guyot A, Clements J, Bradley N (2013) Synovial fluid white cell and differential count in the diagnosis or exclusion of prosthetic joint infection. *Bone Joint J* 95-B: 554–557
9. Kim SG, Kim JG, Jang KM, et al (2017) Diagnostic value of synovial white blood cell count and serum c-reactive protein for acute periprosthetic joint infection after knee arthroplasty. *J Arthroplasty* 32: 3724–3728
10. Ting NT, Della Valle CJ (2017) Diagnosis of periprosthetic joint infection – an algorithm-based approach. *J Arthroplasty* 32: 2047–2050
11. Claassen L, Ettinger S, Pastor MF, et al (2016) The value of arthroscopic neosynovium biopsies to diagnose periprosthetic knee joint low-grade infection. *Arch Orthop Trauma Surg* 136: 1753–1759
12. Klatte TO, Schneider MM, Citak M, et al (2013) Infection rates in patients undergoing primary knee arthroplasty with pre-existing orthopaedic fixation-devices. *Knee* 20: 177–180
13. Szczesny G, Interewicz B, Swoboda-Kopeć E, et al (2008) Bacteriology of callus of closed fractures of tibia and femur. *J Trauma* 65: 837–842
14. Knabl L, Kuppelwieser B, Mayr A, et al (2019) High percentage of microbial colonization of osteosynthesis material in clinically unremarkable patients. *Microbiologyopen* 8: e658
15. Renz N, Feigl S, Dlaska CE, et al (2017) Osteosyntheseassoziierte Infektionen. Epidemiologie, Definition und Diagnostik. *Unfallchirurg* 120: 454–460
16. Trampuz A, Piper KE, Jacobson MJ, et al (2007) Sonication of removed hip and knee prostheses for diagnosis of infection. *N Engl J Med* 357: 654–663
17. Massin P, Bonnin M, Paratte S, et al; French Hip Knee Society (SFHG) (2011) Total knee replacement in post-traumatic arthritic knees with limitation of flexion. *Orthop Traumatol Surg Res* 97: 28–33
18. Rao AJ, Kempton SJ, Erickson BJ, et al (2016) Soft tissue reconstruction and flap coverage for revision total knee arthroplasty. *J Arthroplasty* 31: 1529–1538
19. Tetsworth K, Paley D (1994) Malalignment and degenerative arthropathy. *Orthop Clin North Am* 25: 367–377
20. Lonner JH, Siliski JM, PA Lotke (2000) Simultaneous femoral osteotomy and total knee arthroplasty for treatment of osteoarthritis associated with severe extra-articular deformity. *J Bone Joint Surg Am* 82: 342–348
21. Frosch KH, M Krause M, Frings J, et al (2016) Posttraumatische Deformitäten am Kniegelenk. Intraartikuläre Korrekturosteotomien nach fehlverheilten Tibiakopffrakturen. *Unfallchirurg* 119: 859–876